

# Encorafenib and Binimetinib

## Indication

Unresectable or metastatic BRAF V600 mutation-positive melanoma.

(NICE TA562)

## ICD-10 codes

Codes with a prefix C43

## Regimen details

Day	Drug	Dose	Route
1-28	Encorafenib	450mg* OD	PO
1-28	Binimetinib	45mg BD	PO

**\*Administration of encorafenib at a dose of 450 mg once daily as a single agent is not recommended. If binimetinib is temporarily interrupted, encorafenib should be reduced to 300 mg once daily during the time of binimetinib dose interruption because encorafenib is not well-tolerated at the dose of 450 mg as a single agent**

## Cycle frequency

As above

## Number of cycles

Continuous until disease progression or unacceptable toxicity.

## Administration

Encorafenib is available as 50mg and 75mg capsules. Capsules should be swallowed whole with water and may be taken with or without food. If a dose of encorafenib is missed, the patient should only take the missed dose if it is more than 12 hours until the next scheduled dose.

Grapefruit and grapefruit juice should be **avoided** whilst taking encorafenib.

Binimetinib is available as 15mg tablets. The doses should be taken 12 hours apart. Tablets should be swallowed whole with water and may be taken with or without food. If a dose is missed it should not be taken if it is less than 6 hours until the next dose is due.

If a patient vomits after taking a dose, the dose should not be retaken and the next dose should be taken at the next scheduled time.

## Pre-medication

Nil

## Emetogenicity

This regimen has mild emetic potential.

## Additional supportive medication

Antiemetics if required.

## Extravasation

N/A

## Investigations – pre first cycle

Before commencing treatment BRAF V600 mutation must be confirmed

Investigation	Validity period ((or as per consultant instruction)
FBC	14 days
U+E (including creatinine)	14 days
LFTs	14 days
Magnesium	14 days
Calcium	14 days
LDH	14 days
Pregnancy test (if applicable)	14 days
Blood pressure	Baseline
ECG (QTc < 500ms) and echocardiogram	Baseline
Echocardiogram*	Baseline

\* if urgent treatment is required then this should not be deferred whilst waiting for an echocardiogram if there is no history of or clinical suspicion of heart failure.

Consider dermatological evaluation.

## Investigations – pre subsequent cycles

Investigation	Validity period
FBC	48 hours
U+E (including creatinine)	48 hours
LFTs	48 hours
Magnesium	48 hours
LDH*	48 hours
Blood pressure	Monthly
ECG	should be monitored before treatment, after the first month, then approximately 3 monthly or more frequently if clinically indicated
Echocardiogram	should be monitored before treatment, after the first month, then approximately 3 monthly or more frequently if clinically indicated

\*Haemolysed LDH result should not stop treatment.

## Standard limits for administration to go ahead

If blood results not within range, authorisation to administer **must** be given by prescriber/ consultant

Investigation	Limit
Neutrophils	$\geq 1.0 \times 10^9/L$
Platelets	$\geq 75 \times 10^9/L$
Creatinine clearance (CrCl)	$\geq 30\text{ml/min}$
AST/ALT	$\leq 2.5 \times \text{ULN}$ (or $<5 \times \text{ULN}$ if liver metastases)
Bilirubin	$\leq 1.5 \times \text{ULN}$
QTc	$< 500\text{ms}$ and $<60\text{ms}$ increase from baseline
LVEF	$> \text{LLN}$ for institution

## Dose modifications

Dose modifications should be made as per the table below:

Dose level	Encorafenib dose	Binimetinib dose
Full dose	450mg OD	45mg BD
First reduction	300mg OD	30mg BD
Second reduction	200mg OD	Further dose reductions are not recommended. Discontinue if 30mg BD not tolerated.
Third reduction	100mg OD	

Dose reductions beyond these levels are not recommended.

Administration of encorafenib at a dose of 450 mg once daily as a single agent is not recommended. If binimetinib is temporarily interrupted, encorafenib should be reduced to 300 mg once daily during the time of binimetinib dose interruption because encorafenib is not well-tolerated at the dose of 450 mg as a single agent. If encorafenib is temporarily interrupted (see Table 2), binimetinib should be interrupted. If either agent is permanently discontinued, then both should be discontinued.

If treatment-related toxicities occur, then encorafenib and binimetinib should be simultaneously dose reduced, interrupted or discontinued. Exceptions where dose modifications are necessary for binimetinib only (adverse reactions primarily related to binimetinib) are: retinal pigment epithelial detachment (RPED), retinal vein occlusion (RVO), interstitial lung disease/pneumonitis, cardiac dysfunction, CK elevation and rhabdomyolysis, and venous thromboembolism (VTE).

- **Haematological toxicity**

If neutrophils  $< 1.0 \times 10^9/L$  and/or platelets  $< 75 \times 10^9/L$  consider withholding treatment.

See below for management of pyrexia.

- **Renal impairment**

No dose reduction necessary for mild to moderate renal impairment. Use encorafenib with caution and closely monitor if severe renal impairment.

No dose adjustment of binimetinib is recommended for patients with renal impairment.

- **Hepatic impairment**

No dose modification is required for mild hepatic impairment. Encorafenib and binimetinib are not recommended in moderate or severe hepatic impairment.

- **Other toxicities**

### Pyrexia

Treatment should be interrupted if the patient's temperature is  $\geq 38.5^\circ C$ . Patients should be evaluated for signs and symptoms of infection. Treatment can be restarted once the fever resolves with appropriate prophylaxis using non-steroidal anti-inflammatory medicinal products or paracetamol. If fever is associated with other severe signs or symptoms, treatment should be restarted at a reduced dose once fever resolves and as clinically appropriate.

Toxicity	Grade	Action
Cutaneous	Grade 1-2	Continue If worsens or does not improve within 2 weeks withhold binimetinib and encorafenib until $\leq$ Grade 1 then resume at same dose. If recurs resume with one dose level reduction.
	Grade 3	Withhold binimetinib and encorafenib until $\leq$ Grade 1 then resume at same dose. If recurs resume with one dose level reduction.
	Grade 4	Discontinue
Palmar Plantar Erythrodysesthesia (PPE)	Grade 2	Continue with supportive measures If worsens or does not improve within 2 weeks withhold encorafenib until $\leq$ Grade 1 then resume at full dose or with one dose level reduction.
	Grade 3	Withhold encorafenib and use supportive measures. Assess weekly. When improved to $\leq$ Grade 1 then resume at same dose or with one dose level reduction.
Ocular	Grade 2-3 Symptomatic retinal pigment epithelial detachments (RPED)	Withhold binimetinib for up to 2 weeks with ophthalmic monitoring. <ul style="list-style-type: none"> <li>- If improves to <math>\leq</math> Grade 1 resume at same dose</li> <li>- If improves to <math>\leq</math> Grade 1 resume with one dose level reduction</li> <li>- If does not improve to <math>\leq</math> Grade 2 discontinue.</li> </ul>
	Grade 4 (RPED with reduced visual acuity) Or Retinal vein occlusion (RVO)	Discontinue
	Uveitis	If Grade 1-2 does not respond to topical therapy or if Grade 3 withhold encorafenib and repeat ophthalmic monitoring. If Grade 1 and improves to Grade 0 resume with same dose. If Grade 2-3 and improves to Grade 0 or 1 then resume with one dose level reduction. If not improved in 6 weeks discontinue. Grade 4 – discontinue
Cardiac	Grade 2 LVEF decrease	If asymptomatic withhold binimetinib for up to 4 weeks. It may be recommenced at one dose level reduction if all the following occur within 4 weeks: <ul style="list-style-type: none"> <li>- LVEF <math>\geq</math> LLN</li> <li>- Absolute decrease from baseline is 10% or less.</li> </ul> If LVEF does not recover within 4 weeks – discontinue
	Grade 3-4 LVEF	Discontinue Evaluate LVEF every 2 weeks until recovery
	QTc prolongation $>500$ ms and $\leq 60$ ms from baseline	Withhold encorafenib. Resume with one dose level reduction when $\leq 500$ ms If recurs – discontinue
	QTc prolongation $>500$ ms and $> 60$ ms from baseline	Discontinue
CK elevation	Grade 3 (CK 5-10 x ULN) asymptomatic	Continue binimetinib and ensure adequate hydration

	Grade 4 (CK > 10 x ULN) asymptomatic	Withhold binimetinib until Grade 0 or 1. Ensure adequate hydration.
	Grade 3 or 4 (CK > 5 x ULN) with muscle symptoms or renal impairment	Withhold binimetinib until Grade 0 or 1 If resolved within 4 weeks continue with one dose level reduction If does not recover within 4 weeks – discontinue
VTE	Uncomplicated DVT or < Grade 3 PE	Withhold binimetinib until Grade 0 or 1 If improved continue with one dose level reduction If not improved – discontinue
	Grade 4 PE	Discontinue
LFT abnormalities	Grade 2 AST/ALT 3-5 x ULN	Continue. If no improvement within 2 weeks withhold binimetinib and encorafenib until <3 x ULN or baseline and continue with same dose.
	1 <sup>st</sup> occurrence Grade 3 AST/ALT > 5 x ULN and bilirubin > 2 x ULN	Withhold for up to 4 weeks If improved to grade 0-1 continue binimetinib and encorafenib with one dose level reduction If not improved – discontinue
	1 <sup>st</sup> occurrence Grade 4 AST/ALT > 20 x ULN	Withhold binimetinib and encorafenib for up to 4 weeks If improved to grade 0-1 continue binimetinib and encorafenib with one dose level reduction If not improved – discontinue
	Recurrent Grade 3 AST/ALT > 5 x ULN and bilirubin > 2 x ULN	Discontinue
	Recurrent Grade 4 AST/ALT > 20 x ULN	Discontinue
Interstitial lung disease	Grade 2	Withhold binimetinib for up to 4 weeks If improved to grade 0-1 continue binimetinib with one dose level reduction If not improved – discontinue
	Grade 3-4	Discontinue
Any other adverse reaction	Recurrent or intolerable Grade 2 or 1 <sup>st</sup> occurrence Grade 3	Withhold binimetinib and encorafenib for up to 4 weeks If improved to grade 0-1 continue binimetinib and encorafenib with one dose level reduction If not improved – discontinue
	1 <sup>st</sup> occurrence Grade 4	Withhold binimetinib and encorafenib for up to 4 weeks If improved to grade 0-1 continue binimetinib and encorafenib with one dose level reduction If not improved – discontinue
	Recurrent Grade 3	Discontinue
	Recurrent Grade 4	Discontinue

**Adverse effects** - for full details consult product literature/ reference texts

- **Serious side effects**

Cutaneous squamous cell carcinoma  
QT prolongation  
Haemorrhage  
VTE  
Hypersensitivity reactions  
Ophthalmic reactions, including RVO, RPED, uveitis  
Myelosuppression  
Interstitial lung disease

- **Frequently occurring side effects**

Peripheral neuropathy  
Headache, dizziness  
Pyrexia  
Arthralgia, myalgia  
Photosensitivity  
Rash, pruritus  
Nausea and vomiting  
Diarrhoea  
Alopecia  
Raised LFTs  
Hypertension

- **Other side effects**

**Significant drug interactions** – for full details consult product literature/ reference texts

**Coumarin anticoagulants** (e.g. warfarin): avoid.

**Encorafenib**

**Strong CYP3A4 inhibitors:** Concomitant administration of encorafenib with strong CYP3A4 inhibitors should be avoided due to increased encorafenib exposure and potential increase in toxicity.

**Moderate CYP3A4 inhibitors:** Should be co-administered with caution.

**CYP3A4 inducers:** A reduction in encorafenib exposure is likely and may result in reduced efficacy.

**Transporters:** Potential for encorafenib to inhibit renal transporters OCT2, OAT1, OAT3 and hepatic transporters OATP1B1 and OATP1B3 at clinical concentrations. In addition, encorafenib may inhibit P-gp in the gut and BCRP at the expected clinical concentrations.

**CYP3A4 substrates:** Encorafenib is both an inhibitor and inducer of CYP3A4. Concomitant use with agents that are substrates of CYP3A4 (e.g., hormonal contraceptives) may result in increased toxicity or loss of efficacy of these agents. Agents that are CYP3A4 substrates should be co-administered with caution.

Encorafenib is an inhibitor of UGT1A1. Concomitant agents that are substrates of UGT1A1 may have increased exposure and should be administered with caution.

Please see the SPC for a full list of potential medicinal interactions.

### **Binimetinib**

Binimetinib is primarily metabolised through UGT1A1 mediated glucuronidation. The extent of drug interactions mediated by UGT1A1 is unlikely to be clinically relevant. **UGT1A1 inducers** (such as rifampicin and phenobarbital) and inhibitors (such as indinavir, atazanavir, sorafenib) should be co-administered with caution.

**Inducers of CYP1A2 enzymes** (such as carbamazepine and rifampicin) and **inducers of Pgp transport** (such as Saint John's wort or phenytoin) may decrease binimetinib exposure, which could result in a decrease of efficacy.

Binimetinib is a potential inducer of CYP1A2, and caution should be taken when it is used with sensitive substrates (such as duloxetine or theophylline).

Binimetinib is a weak inhibitor of OAT3, and caution should be taken when it is used with sensitive substrates (such as pravastatin or ciprofloxacin).

### **Additional comments**

Women of child bearing potential must be advised to use adequate barrier contraception throughout treatment.

---

### **References**

- <http://www.swscn.org.uk/guidance-protocols/cancer-protocols/> accessed 9 Jul 2020
- National Institute for Health and Clinical Excellence. NICE Technology Appraisal Guidance 562 via [www.nice.org.uk](http://www.nice.org.uk)
- Summary of Product Characteristics – Encorafenib via [www.medicines.org.uk](http://www.medicines.org.uk)
- Summary of Product Characteristics – Binimetinib via [www.medicines.org.uk](http://www.medicines.org.uk)
- Drummer R., et al. Encorafenib plus Binimetinib versus vemurafenib or Encorafenib in patients with BRAF- mutant melanoma (COLUMBUS): a multicentre, open label, randomised phase 3 trial. *Lancet Oncology*. 2018. 19:5, 603-615.

---

**THIS PROTOCOL HAS BEEN DIRECTED BY DR BOARD, DESIGNATED LEAD CLINICIAN FOR MELANOMA**

**RESPONSIBILITY FOR THIS PROTOCOL LIES WITH THE HEAD OF SERVICE**

Date: Jul 2020

Review: Jul 2023

VERSION: 3

---