

### **Sample Stock list**

Example drugs that a Virtual Ward/Hospital at Home service may wish to consider keeping as part of ward stock/medic's bag.

*Please note: Local formularies may differ. Services should consult with local clinicians, including antimicrobial stewardship leads, when developing their stock lists.*

Drug	Medics bag	In ward stock	Indication
<b>Fluids</b>			
0.9% saline 1L 0.9% saline 500ml (fluid challenge)	1	5	Sepsis Bundle Delirium Infection Dehydration
Plasmalyte 1L	1	5	Hypovolemia
0.45% saline/5% dextrose 1L	1	5	Hypoglycaemic Dehydration
0.9% saline 50ml	2	5	IV Frusemide/mixing of drugs/Intravenous Additives.
Water for Injection 10ml Amp (10)	1	2	Dilution
<b>Glucose</b>			
20% glucose IV (500ml)	1	2	Reversing Hypoglycaemia
Glucagon 1mg - IM	1	1	Reversing Hypoglycaemia
Glucogel - O	1	3	Reversing Hypoglycaemia
<b>Diuretic</b>			
Furosemide (50mg) IV	6	6	Heart Failure (See SOP for IV Furosemide)
<b>Antibiotics IV – requires secondary care senior approval, blood cultures taken prior and sampling from urine/skin.</b>			
<ul style="list-style-type: none"> <li><input type="checkbox"/> Co-amoxiclav 500/100mg powder for solution for injection/infusion</li> <li><input type="checkbox"/> Levofloxacin 5mg/ml solution for infusion</li> <li><input type="checkbox"/> Gentamicin 10mg/ml; 40mg/ml solution for injection/infusion</li> <li><input type="checkbox"/> Teicoplanin 200mg or 400mg powder for solution for injection/infusion or oral solution</li> </ul>	2	2	<b>See Local Formularies re. sensitivities</b>  Pneumonia Sepsis Complex UTIs COPD (sensitivities) Cellulitis
<b>LMWH (different Trusts may use different low molecular weight heparins)</b>			
Dalteparin 10,000units	2	2	Treatment Venous thromboembolism
Dalteparin 5000 units	4	15	Risk assessment prophylaxis Venous thromboembolism – immobile patients
Dalteparin 2500 units	2	2	Risk assessment prophylaxis Venous thromboembolism – immobile patients
<b>Antiemetic</b>			
Ondansetron 8mg IV	1	2	Nausea and vomiting
<b>Emergency drugs</b>			
Anaphylactic Pack: Adrenaline 1:1000 0.5mg – anaphylaxis; Hydrocortisone 100mg	3	2	Resus. Pack
<b>Nebulised medicines</b>			

Salbutamol 2.5mg	20 (1 box)		COPD Exacerbations (Nebuliser supplied through Virtual Ward Team)
Ipratropium 500microgram	20 (1 box)		COPD Exacerbations (Nebuliser supplied through Virtual Ward Team)
<b>Spacer and MDI</b>			
Aerochamber		4	COPD Exacerbation
Salbutamol MDI (SALAMOL)	1	2	COPD Exacerbation
<b>Analgesia</b>			
Paracetamol IV 1g	2		Patients who cannot take oral medication
<b>Oral – Over-labelled Medicines for Emergency Situations.</b> <i>For dosage, please consult local formulary/national guidelines</i>			
AMOXICILLIN 500mg CAPSULES (15 caps)		2	Acute Respiratory Infections (ARI) / Community-acquired pneumonia (CAP)
CLARITHROMYCIN 500mg TABLETS (14)		1	Acute Respiratory Infections (allergy to penicillin/ sensitivity discussion with microbiology)/ Community-acquired pneumonia Cellulitis (penicillin allergy)
DOXYCYCLINE 100mg CAPSULES (8)		2	Acute Respiratory Infections Cellulitis
PREDNISOLONE 5mg TABLETS (28 tabs)		3	Acute Respiratory Exacerbations
CO-AMOXICLAV 500/125 TABLETS (21)		1	Cellulitis (second line after flucloxacillin) Community Acquired Pneumonia
FLUCLOXACILLIN 500mg CAPSULES (28 caps)		1	Cellulitis
NITROFURANTOIN 100mg MR CAPSULES (6 caps) (Ensure eGFR>30)		2	Acute urinary tract infection Females
NITROFURANTOIN 100mg MR CAPSULES (14 caps) (Ensure eGFR>30)		1	Acute urinary tract infection – Males
TRIMETHOPRIM 200mg (6 tabs)		1	Acute urinary tract infection – Females
TRIMETHOPRIM 200mg (14 tabs)		1	Acute urinary tract infection – Males
BISACODYL SUPPOSITORIES 10mg (2) (Stimulant laxative)	1	1	Constipation
SENNA 7.5mg Tablets (10) (Stimulant laxative)	1	1	Constipation
MOVICAL SACHETS (30) (Osmotic laxative)	1	1	Constipation
ASPIRIN DISP 300mg (4)	1	2	Chest Pain
GLYCERAL TRINITRATE SPRAY (1)	1	1	Chest Pain
FRUSEMIDE 40mg (8)	1	2	Heart Failure
DIGOXIN 125 micrograms (8)	1	1	AF/Heart Failure
BISOPROLOL 2.5mg TABLETS (28 tabs)		1	Heart Failure

			Tachycardia Hypertension
EDOXABAN TABLETS 60mg (14 tab) Refer to GP/Cardiology for advice and ongoing care (NB. Reduce dose in renal impairment)		1	Atrial Fibrillation – Prevention of stroke and systemic embolism
EDOXABAN TABLETS 60mg (14 tab)  (NB. Duration to be determined by consultant)		1	Treatment of deep vein thrombosis, treatment of pulmonary embolism and prevention of recurrent deep vein thrombosis and pulmonary embolism (Venous thromboembolism)
EDOXABAN TABLETS 30mg (14 tab)		1	Treatment of deep vein thrombosis, treatment of pulmonary embolism and prevention of recurrent deep vein thrombosis and pulmonary embolism (Venous thromboembolism) in patients with one or more of the following: Moderate or severe renal impairment Low body weight ≤ 60 kg Concomitant use of: ciclosporin, dronedarone, erythromycin, or ketoconazole.
ROTIGOTINE 2mg/24hrs Transdermal Patches (2)	1	1	Parkinson's Disease (when oral not possible)
CYCLIZINE 50mg TABLETS (8)		1	Nausea and vomiting
Paracetamol 1g soluble x4	1	1	Pain management
Paracetamol 500mg tablets (24)	1	2	Pain management
Oramorph 10mg/5ml – 100ml		1	Pain management Breathlessness/agitation (COPD)
<b>Oxygen:</b> To be prescribed in emergency via HOOF form			